

Implications of Gauge Unification for Time Variation of the Fine Structure Constant

Paul Langacker*, Gino Segrè and Matthew J. Strassler

*Department of Physics and Astronomy, University of Pennsylvania
Philadelphia, PA 19104*

ABSTRACT: Unification of the gauge couplings would imply that time variations of the fine structure constant are accompanied by calculable and very significant time variations in the QCD scale parameter Λ_{QCD} . Since Λ_{QCD} is the dominant factor in setting the hadron masses, estimates made by simple variations of the fine structure constant may not provide meaningful limits. There may also be related variations in Yukawa couplings and the electroweak scale. Implications for the 21 cm hyperfine transition, big bang nucleosynthesis, and the triple alpha process are discussed. We find that the first of these already provides strong constraints on the underlying theory. It is emphasized more generally that time (and space) variations of fundamental couplings and their correlations may be a significant probe of ultra-high-energy physics.

*On leave at the School of Natural Sciences, Institute for Advanced Study, Einstein Drive, Princeton, NJ 08540

Slow temporal variations of coupling constants are conceivable in many theories, *e.g.*, those in which their values are related to expectation values of scalar fields [1]. (Naturally, spatial variations may also occur [2].) This possibility has been of considerable interest since first proposed by Dirac [3] over sixty years ago. Webb *et al.* [4] have recently reported that the fine structure constant α displays time variation. The group reports a change $\frac{\Delta\alpha}{\alpha} = -(0.72 \pm 0.18) \times 10^{-5}$ over the redshift range $0.5 < z < 3.5$, where $\Delta\alpha$ is α at large redshift minus α at the present day. In this note we will discuss how this result, and other constraints, fit together if the coupling constants of the strong, weak and electromagnetic interactions are unified.

From a purely observational point of view, all fundamental constants, from the cosmological constant on down, should be viewed as potentially having spatial and time variation of an unknown sort. More interesting than the particular variation of any individual couplings would be the correlations in the variations of *different* couplings. Such correlations potentially will tell us how the variations are being induced, and therefore give insight into the underlying theory. In short, temporal and spatial variations in coupling constants may be a little-recognized probe of ultra-high-energy physics.

Webb *et al.*'s [4] data is based on the absorption spectra of distant quasars, but several constraints exist as well at other redshift values [5]. The best laboratory limit is $|\frac{\Delta\alpha}{\alpha}| < 1.4 \times 10^{-14}$ over a period of 140 days [6]. The best geological limit comes from the naturally occurring nuclear reactor at Oklo in Gabon; recent re-analyses of this data [7], under the assumption that only α varies, and that $\frac{\dot{\alpha}}{\alpha}$ is constant, lead to a limit of $\frac{\dot{\alpha}}{\alpha} = (-0.2 \pm 0.8) \times 10^{-17}/\text{yr}$ over two billion years.

There are other ongoing or planned observations which will provide additional limits. Fluctuations in the cosmic microwave background resulting from changing the ionization history of the universe could lead [8] to a measurement with experimental sensitivity of $|\frac{\Delta\alpha}{\alpha}| < 10^{-2} - 10^{-3}$ at $z \sim 1000$. Nucleosynthesis [9] places bounds on the variations of α with roughly the same order of magnitude, though at a much larger redshift, $z \sim 10^9 - 10^{10}$, again assuming that the only variation is in α .

If one assumes that $\frac{\dot{\alpha}}{\alpha}$ remains approximately constant throughout the history of the universe, the resultant value from the Webb *et al.* [4] data is $\frac{\dot{\alpha}}{\alpha} \sim 10^{-15}/\text{yr}$, which is two orders of magnitude above the Oklo bound although compatible with the other limits. There is, however, no particular reason to assume constant time variation, so until the data suggests that $\frac{\dot{\alpha}}{\alpha}$ is constant, there is no clear contradiction between Oklo and the new observations.

The time-variation of other quantities can also be measured using absorption lines. Cowie and Songaila [10] constrain $X \equiv \alpha^2 g_p m_e / M_p$, where the proton magnetic moment is $eg_p / 2M_p$, from the 21 cm hyperfine line in hydrogen at $z \sim 1.8$. Similarly, Potekhin

et al. [11] limit $Y \equiv M_p/m_e$ from molecular hydrogen clouds at $z = 2.81$:

$$\frac{\Delta X}{X} = (0.7 \pm 1.1) \times 10^{-5}, \quad \frac{\Delta Y}{Y} = (8.3^{+6.6}_{-5.0}) \times 10^{-5}. \quad (1)$$

The main point of this paper¹ is to emphasize that the various fundamental constants are likely to vary simultaneously and in a correlated way that depends on the underlying physics, and that studies of such systems as the Oklo reactor, nucleosynthesis, or molecular/atomic absorption lines should take this into account. We will illustrate this by one especially well-motivated example, i.e., by theories in which the gauge couplings unify at some large scale (consistent with the data [13]). Then a change in α will be accompanied by a much larger change in the strong interaction coupling and mass scale. For definiteness, we consider a grand unified gauge theory such as $SU(5)$ with a single coupling constant α_G . Similar considerations would apply to many string theories which compactify directly to the standard model.² Unified theories involving more complicated breaking patterns, exotic matter, higher Kač-Moody levels, or large extra dimensions would differ in the details but not in the qualitative features.

Unification presumably occurs at some large scale M_G , typically of order 3×10^{16} GeV. At lower scales the group breaks down to $SU(3) \times SU(2) \times U(1)$ characterized by couplings α_i , $i = 1, 2, 3$, which however tend to α_G as the scale at which they are measured nears M_G . For instance, in the MSSM (Minimal Supersymmetric Standard Model), the couplings run so that at the electroweak scale M_Z they are given by

$$\alpha_i^{-1}(M_Z) = \alpha_G^{-1} + b_i t_G \quad (2)$$

where $t_G = \frac{1}{2\pi} \ln \frac{M_G}{M_Z} \approx 5.32$ and $\alpha_G^{-1} \approx 23.3$. The b_i are fixed by the relative particle multiplets; in the MSSM one has $b_i = (\frac{33}{5}, 1, -3)$. Between M_G and M_Z the α_i rather than α are the relevant couplings. Below M_Z the standard model group $SU(3) \times SU(2) \times U(1)$ is further broken to $SU(3) \times U(1)_{ELM}$, where $U(1)_{ELM}$ is the electromagnetic gauge group. This leads to a value of the electromagnetic coupling at scale M_Z of

$$\alpha^{-1}(M_Z) = \frac{5}{3} \alpha_1^{-1}(M_Z) + \alpha_2^{-1}(M_Z) = 127.9. \quad (3)$$

Now imagine that α_G^{-1} is the vacuum expectation value of some field $\phi(t)$ which is a slowly varying function of time. (For example, $\phi(t)$ might parametrize the change

¹Some of the issues in this paper have been considered in somewhat similar contexts, for example, in [12]

²There is an order of magnitude discrepancy between the observed and expected unification scales in the simplest such theories, but that is only a 10% effect in $\log M_G/M_Z$. This could be accounted for by threshold or other effects, and would not much affect our conclusions.

in the volume of some compact extra dimensions.) This in turn will induce correlated variation, at any given energy scale below M_G , in the $SU(3) \times SU(2) \times U(1)$ coupling constants. To find the relationship between the couplings, we make a second assumption (which we will relax later) that the changes in α_i^{-1} are dominated by the changes in α_G^{-1} — in other words, that corrections to t_G and to threshold effects, etc., are relatively unimportant.³ We also neglect higher order corrections and assume (correctly, in the majority of models) that $\Delta\alpha(M_Z)/\alpha(M_Z) \simeq \Delta\alpha/\alpha$, where $\alpha \equiv \alpha(0) \sim 1/137$ is the conventional low-energy fine structure constant. We then have

$$\frac{\Delta\alpha_i}{\alpha_i} = -\frac{\Delta\alpha_i^{-1}}{\alpha_i^{-1}} = -\frac{\Delta\alpha_G^{-1}}{\alpha_i^{-1}}, \quad (4)$$

so that at scale M_Z

$$\frac{\Delta\alpha}{\alpha} = -\frac{8}{3} \frac{\Delta\alpha_G^{-1}}{\alpha^{-1}} = \frac{8}{3} \frac{\alpha}{\alpha_G} \frac{\Delta\alpha_G}{\alpha_G} \sim 0.49 \frac{\Delta\alpha_G}{\alpha_G}. \quad (5)$$

Similarly, the $SU(3)$ group's coupling, usually called α_S instead of α_3 , satisfies, again at scale M_Z ,

$$\frac{\Delta\alpha_S}{\alpha_S} \simeq \frac{3}{8} \frac{\alpha_S}{\alpha} \frac{\Delta\alpha}{\alpha} \sim 5.8 \frac{\Delta\alpha}{\alpha} \quad (6)$$

i.e., the proportional variation in the strong subgroup's coupling is almost six times as great as that of α , when measured at the scale M_Z , where $\frac{\alpha_S}{\alpha} \simeq 15.4$.

The physical quantities we are most directly interested in are the associated relative changes in the neutron and proton masses and, to the extent this is possible to estimate, the changes in nuclear binding energies. The nucleon masses are largely fixed by the QCD scale Λ_{QCD} . The expression we are most interested in is therefore the relation of $\frac{\Delta\Lambda_{QCD}}{\Lambda_{QCD}}$ to $\frac{\Delta\alpha}{\alpha}$ (up and down quark masses make only a minute contribution to nucleon masses). Λ_{QCD} is approximately the scale at which α_S diverges, $\alpha_S^{-1}(\Lambda_{QCD}) = 0$. To see how changes in this scale are related to changes in α , we must study the further running of α_S from the scale M_Z down to μ . This change is again given by the renormalization group equations

$$\alpha_S^{-1}(\mu) = \alpha_S^{-1}(M_Z) - \frac{b_3^{\text{SM}}}{2\pi} \ln \left(\frac{\mu}{M_Z} \right) \quad (7)$$

where $\alpha_S^{-1}(M_Z) = 8.33$ and $b_3^{\text{SM}} = -11 + \frac{2}{3}n_F$, where n_F is the number of quark flavors light compared to μ .

³Clearly we are also assuming that the back-reaction of a varying α_G on the cosmological constant is not large; we will comment on this below.

Using the above equation, the value of μ at which $\alpha_S^{-1}(\mu) \rightarrow 0$ is given by

$$\Lambda_{QCD}^9 \sim M_Z^{23/3} m_b^{2/3} m_c^{2/3} \exp\left(-\frac{2\pi}{\alpha_S(M_Z)}\right). \quad (8)$$

where m_b, m_c are the bottom and charm quark masses, whose subleading dependence on α_S we are ignoring. This leads immediately to the desired relation

$$\frac{\Delta\Lambda_{QCD}}{\Lambda_{QCD}} = -\frac{2\pi}{9} \Delta\alpha_S^{-1}(M_Z) = \frac{2\pi}{9\alpha_S} \frac{\Delta\alpha_S}{\alpha_S} \quad (9)$$

where α_S is measured at M_Z . Using (6), we finally arrive at a prediction

$$\frac{\Delta\Lambda_{QCD}}{\Lambda_{QCD}} \sim 34 \frac{\Delta\alpha(M_Z)}{\alpha(M_Z)} \sim 34 \frac{\Delta\alpha}{\alpha}. \quad (10)$$

(The coefficient has a theoretical uncertainty at the level of perhaps twenty percent.) This suggests that, within the framework of a unified theory and assuming the nucleon mass scale is set by Λ_{QCD} , a variation such as the one suggested by Webb *et al.* [4] of $\frac{\Delta\alpha}{\alpha} = -0.72 \times 10^{-5}$ should be accompanied by a shift in $\frac{\Delta M_p}{M_p} \sim -25 \times 10^{-5}$, clearly a very significant correction.

Before discussing the implications, let us consider possible variations in other physical constants. These are likely to occur along with those in the gauge couplings, although the specific relation cannot be obtained without a more specific theory. However, it is useful to parametrize them and consider likely possibilities.

Only dimensionless quantities such as coupling constants or ratios of masses are physically significant. We take the point of view that the unification scale M_G is simply a reference scale relative to which other masses such as Λ_{QCD} are expressed. Indeed, the formula (10) should more properly be interpreted, in this language, as a statement about the variation in the quantity Λ_{QCD}/M_G . Later we will have to consider the variation in the Newton constant $G_N \propto M_{pl}^{-2}$ relative to M_G .

We next consider the (running) Yukawa coupling h_a of fermion a , which is related to its mass by $m_a = h_a v$, where $v \sim 246$ GeV is the electroweak scale. (For the quarks, the physical mass is $\sim (m_q + M_{QCD})$, where $M_{QCD} \sim \Lambda_{QCD}$, the latter contribution dominating for u and d .) We expect that since the gauge coupling is varying at the unification scale, the same will be true, to some degree, for the Yukawa couplings. Since the size of this effect is model-dependent, we parametrize our ignorance by introducing unknown constants λ_a , where

$$\frac{\Delta h_a(M_G)}{h_a(M_G)} \equiv \frac{\lambda_a}{2} \frac{\Delta\alpha_G}{\alpha_G} \sim \lambda_a \frac{\Delta\alpha}{\alpha}. \quad (11)$$

The running of h_a is calculable in a given model. For the light quarks and leptons, the running is dominated by the gauge contributions, yielding

$$\frac{h_a(M_G)}{h_a(M_Z)} = \prod_{i=1,2,3} \left[\frac{\alpha_G(M_G)}{\alpha_i(M_Z)} \right]^{\frac{b_{a;i}}{2b_i}} \quad (12)$$

where in the MSSM the anomalous dimensions are given by $b_{u;i} = \left(\frac{-13}{15}, -3, \frac{-16}{3}\right)$, $b_{d;i} = \left(\frac{-7}{15}, -3, \frac{-16}{3}\right)$ and $b_{e;i} = \left(\frac{-9}{5}, -3, 0\right)$. This implies

$$\frac{\Delta h_a(M_Z)}{h_a(M_Z)} \sim \left(\lambda_a - \sum_i b_{a;i} \alpha_i t_G \right) \frac{\Delta \alpha}{\alpha}, \quad (13)$$

where α_i is evaluated at M_Z . One finds $-\sum_i b_{a;i} \alpha_i t_G \sim 4.0$ for $a = u$ or d , and 0.70 for $a = e$. That is, there is a moderate magnification of the quark Yukawas from the gauge corrections (mainly from α_S), but a much smaller effect for the electron. One might expect that this effect for the quarks gets further magnified as one runs from M_Z down to the quark mass. This is true for the bottom and charm Yukawa couplings. However, for the light quarks, the running should be taken down to a scale of order Λ_{QCD} . When this is done, and the variation of Λ_{QCD} is accounted for, it turns out there is a cancellation and the effect is actually smaller. Given the inherent theoretical uncertainties in our computations, we have chosen to ignore this small shift, and will simply use the λ_a for the light quarks.

More important are the changes in the electroweak scale v , which are tied to the scale of supersymmetry breaking in most supersymmetric models. For example, in the traditional supergravity mediated models there are various soft supersymmetry breaking masses (scalar masses, sfermion masses, and other bilinear and cubic scalar terms), which are usually assumed to have the same order of magnitude m_{soft} at the Planck scale.⁴ Although these soft parameters run (and may change sign for scalar mass-squares), they are generally of the same order of magnitude at the weak scale. Typically, one finds that v^2 (and the inverse G_F^{-1} of the Fermi constant) scales as m_{soft}^2/α_{weak} , where $\alpha_{weak} = \frac{3}{5}\alpha_1 + \alpha_2$, and that the corresponding W and Z masses scale as m_{soft} . The underlying mechanism for breaking supersymmetry in the hidden sector and therefore generating m_{soft} is unknown. We shall parametrize our ignorance by introducing

$$\frac{\Delta v}{v} \equiv \kappa \frac{\Delta \alpha}{\alpha}. \quad (14)$$

⁴We are assuming that μ , the supersymmetric Higgs mass, is comparable to the soft parameters, whether it is elementary or dynamically generated.

Then,

$$\frac{\Delta m_a}{m_a} \sim (\lambda_a - \sum_i b_{a;i} \alpha_i t_G + \kappa) \frac{\Delta \alpha}{\alpha}. \quad (15)$$

In many supersymmetry-breaking models, a dynamical strong-coupling scale sets an intermediate supersymmetry breaking scale M_I , which in turn feeds into m_{soft} . Typically this relation takes a form $m_{soft} \sim M_I^n / M_G^{n-1}$, n positive. It is easy to see that if the supersymmetry-breaking dynamics is also unified with the standard model, with coupling constant α_G at M_G , then for any n

$$\frac{\Delta m_{soft}}{m_{soft}} \sim -\frac{\Delta \alpha_G}{\alpha_G} \ln \frac{m_{soft}}{M_G} \sim 35 \frac{\Delta \alpha_G}{\alpha_G} \sim 70 \frac{\Delta \alpha}{\alpha} \quad (16)$$

or, in short, $\kappa \sim 70$. (Note that there is no corresponding reason to expect the λ_a to be large.) However, the coupling constant of the supersymmetry-breaking physics may be set in other ways, in which case κ is completely unconstrained.

One can now go back and investigate the correctness of our original assumption of considering only the change in α_G^{-1} and not that in t_G in (2). It is easily shown that the effect of the change in t_G induced by Δv is not small when $|\kappa|$ is of order 10 or larger. For example, the second and third expressions in (6) acquire correction factors $(1 - 10 \frac{\alpha}{\pi} \kappa) \sim (1 - 0.025 \kappa)$. It is appropriate, then, to recalculate the variation of Λ_{QCD} to see whether we have left out a large effect. We find

$$\frac{\Delta \Lambda_{QCD}}{\Lambda_{QCD}} \sim 34(1 + .005 \kappa) \frac{\Delta \alpha}{\alpha}. \quad (17)$$

where the coefficient of κ is uncertain at the level of twenty percent due to ambiguities in the one-loop renormalization prescription. Note that the effect of $\kappa \sim 70$ is substantial but not overwhelming. The uncertainties in its effect are comparable to or below the errors in [4], as we will see.

We finally mention that changes in the gauge couplings may also be correlated with changes in the cosmological constant Λ . Indeed, it has recently been argued that fine-tuned cancellations involving radiative contributions would be upset by a time variation in α , with enormous effect [14]. Since the smallness of Λ is the outstanding puzzle in particle physics, it is difficult to know how it is affected. Naively, quantum field theory and classical gravity would suggest the back-reaction due to varying couplings would be enormous, but we already know that effective field theory reasoning is wrong, and do not know why. Since there is neither theoretical nor experimental guidance on this point, we choose to assume the cosmological constant is not affected; whether this is correct is, in our view, still an experimental question.

As a first application, let us consider the quantities $X \equiv \alpha^2 g_p m_e / M_p$ and $Y \equiv M_p / m_e$. For the nucleon masses, one has

$$\begin{aligned} M_p &= M_{nuc} + 2m_u + m_d + \alpha M_{elm} \\ M_n &= M_{nuc} + m_u + 2m_d, \end{aligned} \quad (18)$$

where $M_{nuc} \sim 3M_{QCD}$ scales like Λ_{QCD} . αM_{elm} is the electromagnetic contribution to M_p , with M_{elm} also scaling like Λ_{QCD} . The variations of $m_{u,d}$ and m_e are given in (15). $m_{u,d}$ are typically estimated to be a few MeV, so that $M_{p,n}$ are dominated by M_{nuc} . Hence, we expect

$$\frac{\Delta M_p}{M_p} \sim \frac{\Delta \Lambda_{QCD}}{\Lambda_{QCD}}. \quad (19)$$

(This would have to be modified if λ_a or κ were enormous.) The proton and neutron g factors $g_{p,n}$ are well described in the constituent quark model, where they are Clebsch-Gordan coefficients, so we will ignore possible variation in g_p . Assuming that $\lambda_u \approx \lambda_d \approx \lambda_e$, and denoting their average value as λ , one then has

$$\begin{aligned} \frac{\Delta X}{X} &\sim (-32 + \lambda + 0.8\kappa) \frac{\Delta\alpha}{\alpha} \sim (23 \pm 6) \times 10^{-5} \\ \frac{\Delta Y}{Y} &\sim (34 - \lambda - 0.8\kappa) \frac{\Delta\alpha}{\alpha} \sim (-24 \pm 6) \times 10^{-5}, \end{aligned} \quad (20)$$

where the numerical values are obtained using the Webb *et al.* value $\frac{\Delta\alpha}{\alpha} = (-0.72 \pm 0.18) \times 10^{-5}$ and $\lambda = \kappa = 0$. (The theoretical errors in this computation, which are slightly smaller than the experimental errors but could straightforwardly be reduced by more careful calculation, are not shown.) One sees the large magnification implied by coupling constant unification. The values in (20) are to be compared with the observational limits [10, 11], given in eq. (1), obtained at redshifts within the Webb *et al.* range. Consistency of (20) and (1) would require a rather delicate cancellation, with $\lambda + 0.8\kappa \sim 32$.

We should emphasize that one ought not to conclude from this that the result of [4] is inconsistent with unification of coupling constants. Even if $\lambda + 0.8\kappa$ lies outside the range 25 – 35, one can conclude only that unification requires that the variation observed by Webb *et al.* is caused by physics at distance scales long compared with M_G^{-1} , and preferentially affecting electromagnetic or electroweak physics. Our aim is to use [4] to constrain the theory, not the reverse.

Let us now turn to another process which also could be strongly affected by coupling variations: nucleosynthesis. Consider the neutron-proton ratio in the early universe, fixed approximately at the temperature T_F at which the the weak-interaction processes

that interconvert neutrons and protons freeze out by comparison to the expansion rate. The relative abundance is

$$\frac{N_n}{N_p} \simeq e^{\frac{-(M_n - M_p)}{T_F}}, \quad (21)$$

where by (18) the neutron-proton mass difference $M_n - M_p \sim 1.29$ MeV is given by

$$M_n - M_p = m_d - m_u - \alpha M_{elm}. \quad (22)$$

For definiteness, we will use the estimates [15] $m_d - m_u = 2.05 \pm 0.30$ MeV, and $\alpha M_{elm} = 0.76 \pm 0.30$ MeV. The freeze-out temperature $T_F \sim 0.72$ MeV is set roughly by equating the weak interaction rate at temperature T to the universe's expansion rate. Expressing the Fermi constant G_F in terms of the Higgs vacuum expectation value v gives

$$\frac{T^5}{v^4} \simeq K \frac{T^2}{M_{Pl}} \implies T_F \simeq \left(K \frac{v^4}{M_{Pl}} \right)^{\frac{1}{3}}, \quad (23)$$

where K is a constant. Using (14) and defining

$$\frac{\Delta M_{pl}}{M_{pl}} = \rho \frac{\Delta \alpha}{\alpha} \quad (24)$$

(recall we are holding M_G fixed,) we have

$$\frac{\Delta T_F}{T_F} = \left(\frac{4}{3}\kappa - \frac{1}{3}\rho \right) \frac{\Delta \alpha}{\alpha}. \quad (25)$$

The variation in $r \equiv N_n/N_p$ is given by

$$\frac{\Delta r}{r} = -\Delta \left(\frac{M_n - M_p}{T_F} \right). \quad (26)$$

In addition to the explicit α dependence, there may be a much larger variation due to M_{elm} , and possibly the quark mass difference and T_F . From (15), (18), and (25) we find

$$\frac{\Delta r}{r} \sim 37(1 - 0.08\lambda - 0.01\kappa - 0.02\rho) \left(\frac{\Delta \alpha}{\alpha} \right)_{\text{BBN}}, \quad (27)$$

where the subscript BBN indicates that the quantity is to be evaluated at the time of nucleosynthesis. Possible changes in r can be estimated from the ${}^4\text{He}$ abundance [16], with typical estimates implying

$$-0.1 < \frac{\Delta r}{r} < 0.02, \quad (28)$$

implying

$$-0.003 < (1 - 0.08\lambda - 0.01\kappa - 0.02\rho) \left(\frac{\Delta\alpha}{\alpha} \right)_{\text{BBN}} < 0.0005. \quad (29)$$

(Note that the coefficient could be negative for $\lambda + 0.8\kappa \sim 32$.) If we assume a constant $\dot{\alpha}/\alpha$, this implies the range $(-3 \times 10^{-13}/\text{yr}, 5 \times 10^{-14}/\text{yr})$ for $(1 - 0.08\lambda - 0.01\kappa - 0.02\rho) \frac{\dot{\alpha}}{\alpha}$, which is consistent with the result of [4]. However, other time dependences (e.g., a linear dependence on $1 + z$) could give very different results; and we already know that, for constant $\frac{\dot{\alpha}}{\alpha}$, Oklo and the results of [4] are inconsistent (but see below.) Thus it is vital to measure the time dependence of α at moderate z with high precision.

Leaving aside time variation for the moment, we may recall that the actual values of fundamental couplings have themselves been the subject of continued interest, particularly in light of the fact that a universe like ours may only be possible for a narrow range of parameters. This subject, commonly known by the catchword of the “anthropic principle” [17], bears on the issue of varying couplings. If the time dependence of the couplings was such as to bring them outside the range of allowed values for certain reactions, the anthropic principle applied at an earlier era might forbid an otherwise acceptable time variation.

In particular, there are contexts in nuclear physics where high sensitivity to varying coupling constants has been noted. One might wonder, then, whether the observed variation might in turn affect the observed properties of stars, or affect the natural reactor at Oklo sufficiently to warrant a reanalysis of results obtained from it.

Agrawal *et al.* [18] have discussed at length, in a variety of settings, the anthropic principle’s constraints on the value of v , the Higgs vacuum expectation value. Recent extensions of this analysis have focused on the much tighter constraints placed by the existence of the triple-alpha process ${}^4\text{He} + {}^8\text{Be} \longrightarrow {}^{12}\text{C}^*$ where ${}^{12}\text{C}^*$ is an excited state of carbon, 7.6 MeV above the ground state. The existence of this resonance and the non-existence of a resonance below threshold in oxygen ensures that (1) carbon is produced in stellar interiors and (2) the carbon is not immediately converted into oxygen. In other words the existence of carbon in our universe is sensitively dependent on the location of nuclear resonances in both carbon and in oxygen. The subject becomes particularly interesting for our considerations because Livio *et al.* [19] and more recently Oberhummer *et al.* [20] have shown how sensitively the energies of these resonances depends on the basic structure of the nucleon-nucleon potential. As an illustration, consider carbon production in stars, which has been occurring in stars for many billions of years: the time variation of α cannot be such as to imply a change in the nucleon-nucleon potential by a quantity sufficient to shut down carbon production. There is some question of how much of a change in the nucleon-nucleon potential is acceptable, but it is apparently of order 1%.

A shift in Λ_{QCD} , as in equation (10), is by itself not very important, because the main features of nuclei (nucleon masses, nuclear potential depths) shift along with it. At leading order, Λ_{QCD} is the only scale affecting QCD physics, so that all dimensionless quantities are shift-invariant. However, this scaling is broken by the quark masses; and in stars, it is also broken by any associated changes in the stellar environment, such as the core temperature, which might be induced by changes in the various parameters (including $M_{pl.}$). In particular, unlike most hadronic masses, the pion mass does not scale like Λ_{QCD} . This is because of its special role as a pseudo-Goldstone boson of an approximate global chiral $SU(2)$ symmetry of the strong interactions, which is explicitly broken by the small quark mass terms. One has $m_\pi^2 \sim f_\pi (m_u + m_d)$ with the pion decay constant $f_\pi \sim \Lambda_{QCD}$. At some level this effect will feed into other hadronic masses, such as that of the ρ , and modify their linear scaling, and will also affect such dimensionless quantities as the pion-nucleon coupling. However, the dominant effect should be on the range of the pion exchange potential, which effectively shifts the strength by a factor of $-\Delta(m_\pi/M_{had})$, where M_{had} is a typical hadronic scale. If we assume for illustration that this must be $< 1\%$, one obtains

$$\left| \frac{\Delta(m_\pi/M_{had})}{(m_\pi/M_{had})} \right| \sim \frac{1}{2} \left| \frac{\Delta(m_q/\Lambda_{QCD})}{m_q/\Lambda_{QCD}} \right| \sim \frac{1}{2} \left| (-34 + \lambda + 0.8\kappa) \frac{\Delta\alpha}{\alpha} \right| < 0.01. \quad (30)$$

The limit on $\frac{\Delta\alpha}{\alpha}$ is then of order $10^{-2} - 10^{-3}$ over a period of five billion years, a constraint certainly compatible with the Webb *et al.* [4] observation. (This simple estimate is consistent, when appropriately translated, with the more thorough calculation of Jeltama and Sher [21], who considered the changes induced in the potential by variations in v , keeping $\alpha_S(M_Z)$ fixed.) A full calculation of the effect of varying m_π/M_p , using a realistic nuclear potential model along the lines of [21], and taking into account possible variations in the stellar environment, would be useful.

Meanwhile, the Oklo reactor data have so far been analyzed in detail only for the effects of a direct variation of α . Changes in the effective nuclear potential, especially from the change in m_π/M_{had} , could have a significant effect on our understanding of its implications. In particular, the statement that the combined results of Webb *et al.* and those of Oklo are inconsistent with constant $\frac{\dot{\alpha}}{\alpha}$ assumes that only α is varying; if the nuclear physics is also varying, then no firm conclusion is possible at this time. A reanalysis of the Oklo reactor, aimed at obtaining more accurately the constraints that it imposes on the space of coupling constants, would therefore be welcome. It is possible that this reanalysis would put strong constraints on the variation of Λ_{QCD}/v , as well as on α , and these would be interesting to know.

To summarize, we have considered the possibility that the variation of coupling constants is caused by physics at very short distance scales, where the gauge couplings

may be unified. If this is the case, one should not treat the variation of the fine structure constant in isolation. One must simultaneously treat the variations in the strong coupling; the running of coupling constants suggest these are very significant. Within our framework a measured variation of an expression of the form $\alpha \frac{m_e}{M_p}$, for example, is due more to changes of $\frac{m_e}{M_p}$ than of α . We have also calculated the changes induced in Yukawa couplings by anomalous dimensions. Parametrizing the changes in Yukawa couplings at the unification scale by λ and in the electroweak scale v by κ , respectively, we find the Webb *et al.* observations are consistent with other absorption line results (which depend on m_e/M_p) only for specific and large values for $\lambda + 0.8\kappa$. Large effects are also possible for the abundance of primordial ${}^4\text{He}$, but these are at a much earlier period and cannot be directly related to the absorption line results without a detailed model of the time dependence. A reanalysis of the Oklo reactor constraints and a detailed study of the triple alpha process, taking into account these types of effects, would be extremely useful.

More generally, the study of temporal and spatial variations in fundamental constants is potentially a powerful probe of fundamental physics. Both the time-variation of these couplings (which can be constrained by combining, *e.g.*, nucleosynthesis, cosmic microwave radiation, and absorption line data), and the correlations amongst any observed variations at a fixed time, provide strong diagnostics of the underlying theoretical structure. It is important that thorough and dedicated efforts to constrain such variation be made.

As this work was completed, a paper by Calmet and Fritzsche appeared [22] with considerable overlap. Though there are differences in the details, the two papers agree on the main point: that coupling constant unification implies that a time variation in α be accompanied by a much larger variation in strong interaction parameters such as the nucleon mass, and that limits on time variation need to be recalculated and re-interpreted.

All three authors wish to express their gratitude to the Department of Energy for support through grant DOE-EY-76-02-3071. One of us (PL) was also supported by the W. M. Keck Foundation as a Keck Distinguished Visiting Professor at the Institute for Advanced Study and by the Monell Foundation. PL thanks J. Bahcall, F. Dyson and N. Seiberg, and MJS thanks J. Erler, A. Naqvi and G. Shiu, for useful discussions.

References

- [1] See, for example, J. D. Bekenstein, *Phys. Rev. D* **25**, 1527 (1982); T. Damour and A. M. Polyakov, *Nucl. Phys. B* **423**, 532 (1994); B. A. Campbell and K. A. Olive,

- Phys. Lett. B **345**, 429 (1995); E. Witten, hep-ph/0002297; K. A. Olive and M. Pospelov, hep-ph/0110377.
- [2] J. D. Barrow and C. O’Toole, astro-ph/9904116.
 - [3] P. A. M. Dirac, Nature **139**, 323 (1937).
 - [4] J. K. Webb *et al.*, Phys. Rev. Lett. **87**, 091301 (2001).
 - [5] For a general review of constraints on the variation of fundamental constants, see T. Chiba, gr-qc/0110118.
 - [6] J. D. Prestage, R. L. Tjoelker and L. Maleki, Phys. Rev. Lett. **74**, 3511 (1995).
 - [7] T. Damour and F. Dyson, Nucl. Phys. B **480**, 37 (1996); Y. Fujii *et al.*, Nucl. Phys. B **573**, 377 (2000).
 - [8] S. Hannestad, Phys. Rev. D **60**, 023515 (1999); M. Kaplinghat, R. J. Scherrer and M. S. Turner, Phys. Rev. D **60**, 023516 (1999).
 - [9] E. W. Kolb, M. J. Perry and T. P. Walker, Phys. Rev. D **33**, 869 (1986); L. Bergstrom, S. Iguri and H. Rubinstein, Phys. Rev. D **60**, 045005 (1999); P. P. Avelino *et al.*, Phys. Rev. D **64**, 103505 (2001).
 - [10] L. L. Cowie and A. Songaila, Astrophys. J. **453**, 596 (1995).
 - [11] A. Y. Potekhin *et al.*, Astrophys. J. **505**, 523 (1998).
 - [12] W. J. Marciano, Phys. Rev. Lett. **52**, 489 (1984); C. T. Hill and G. G. Ross, Nucl. Phys. B **311**, 253 (1988); C. T. Hill, P. J. Steinhardt and M. S. Turner, Phys. Lett. B **252**, 343 (1990).
 - [13] See, for example, P. Langacker and N. Polonsky, Phys. Rev. D **52**, 3081 (1995).
 - [14] T. Banks, M. Dine and M. R. Douglas, hep-ph/0112059.
 - [15] J. Gasser and H. Leutwyler, Phys. Rept. **87**, 77 (1982).
 - [16] For a recent review, see G. Steigman, astro-ph/0009506.
 - [17] For recent reviews, see C. J. Hogan, Rev. Mod. Phys. **72**, 1149 (2000); B. Müller, astro-ph/0108259.
 - [18] V. Agrawal, S. M. Barr, J. F. Donoghue and D. Seckel, Phys. Rev. Lett. **80**, 1822 (1998) Phys. Rev. D **57**, 5480 (1998).
 - [19] M. Livio, D. Hollowell, A. Weiss and J. W. Truran, Nature **340**, 281 (1989).

- [20] A. Csoto, H. Oberhummer and H. Schlattl, Nucl. Phys. A **688**, 560 (2001);
H. Oberhummer, R. Pichler and A. Csoto, nucl-th/9810057.
- [21] T. Jeltema and M. Sher, Phys. Rev D **61**, 017301 (2000).
- [22] X. Calmet and H. Fritzsch, hep-ph/0112110.